

Black Holes in Nuclear and Particle Physics

Daniel Grumiller

Center for Theoretical Physics
Massachusetts Institute of Technology
and
Institute for Theoretical Physics
Vienna University of Technology

Annual meeting of the Austrian Physical Society, Aflenz, Austria,
September 2008

Outline

Black Holes in Gravitational Physics

Black Holes in Non-Gravitational Physics

Case Study: Holographic Renormalization

Outline

Black Holes in Gravitational Physics

Black Holes in Non-Gravitational Physics

Case Study: Holographic Renormalization

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."
- ▶ A. Eddington (1935): "I think there should be a law of Nature to prevent a star from behaving in this absurd way!"

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."
- ▶ A. Eddington (1935): "I think there should be a law of Nature to prevent a star from behaving in this absurd way!"
- ▶ S. Hawking (1975): "If Black Holes do exist Kip (Thorne) will get one year of Penthouse."

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."
- ▶ A. Eddington (1935): "I think there should be a law of Nature to prevent a star from behaving in this absurd way!"
- ▶ S. Hawking (1975): "If Black Holes do exist Kip (Thorne) will get one year of Penthouse."
- ▶ M. Veltman (1994): "Black holes are probably nothing else but commercially viable figments of the imagination."

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."
- ▶ A. Eddington (1935): "I think there should be a law of Nature to prevent a star from behaving in this absurd way!"
- ▶ S. Hawking (1975): "If Black Holes do exist Kip (Thorne) will get one year of Penthouse."
- ▶ M. Veltman (1994): "Black holes are probably nothing else but commercially viable figments of the imagination."
- ▶ G. 't Hooft (2004): "It is however easy to see that such a position is untenable. (*comment on Veltman*)"

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."
- ▶ A. Eddington (1935): "I think there should be a law of Nature to prevent a star from behaving in this absurd way!"
- ▶ S. Hawking (1975): "If Black Holes do exist Kip (Thorne) will get one year of Penthouse."
- ▶ M. Veltman (1994): "Black holes are probably nothing else but commercially viable figments of the imagination."
- ▶ G. 't Hooft (2004): "It is however easy to see that such a position is untenable. (*comment on Veltman*)"
- ▶ S. Hughes (2008): "Unambiguous observational evidence for the existence of black holes has not yet been established."

Black Holes: A Brief History of Quotes

- ▶ J. Michell (1783): "... all light emitted from such a body would be made to return towards it by its own proper gravity."
- ▶ A. Eddington (1935): "I think there should be a law of Nature to prevent a star from behaving in this absurd way!"
- ▶ S. Hawking (1975): "If Black Holes do exist Kip (Thorne) will get one year of Penthouse."
- ▶ M. Veltman (1994): "Black holes are probably nothing else but commercially viable figments of the imagination."
- ▶ G. 't Hooft (2004): "It is however easy to see that such a position is untenable. (*comment on Veltman*)"
- ▶ S. Hughes (2008): "Unambiguous observational evidence for the existence of black holes has not yet been established."
- ▶ S. Hughes (2008): "Most physicists and astrophysicists accept the hypothesis that the most massive, compact objects seen in many astrophysical systems are described by the black hole solutions of general relativity."

Why Study Black Holes?

Depending whom you ask you'll hear:

- ▶ Mathematician: because they are interesting
- ▶ String Theoretician: because they hold the key to quantum gravity
- ▶ General Relativist: because they are unavoidable
- ▶ Particle Speculator: because they might be produced at LHC
- ▶ Nuclear Physicist: because they are dual to a strongly coupled plasma
- ▶ Astrophysicist: because they explain the data
- ▶ **Cosmologist: because they exist**

But...

Why Study Black Holes?

Depending whom you ask you'll hear:

- ▶ Mathematician: because they are interesting
- ▶ String Theoretician: because they hold the key to quantum gravity
- ▶ General Relativist: because they are unavoidable
- ▶ Particle Speculator: because they might be produced at LHC
- ▶ Nuclear Physicist: because they are dual to a strongly coupled plasma
- ▶ Astrophysicist: because they explain the data
- ▶ Cosmologist: because they exist

But...

Do they exist?

Why Study Black Holes?

Depending whom you ask you'll hear:

- ▶ Mathematician: because they are interesting
- ▶ String Theoretician: because they hold the key to quantum gravity
- ▶ General Relativist: because they are unavoidable
- ▶ Particle Speculator: because they might be produced at LHC
- ▶ Nuclear Physicist: because they are dual to a strongly coupled plasma
- ▶ Astrophysicist: because they explain the data
- ▶ Cosmologist: because they exist


But...

Do they exist?

Let me answer this without getting philosophical, by appealing to data

Some Black Hole Observational Data

OJ287, about 18 billion solar masses




Artistic impression (NASA Outreach), presented at the Annual meeting of the American Astronomical Society, 2008

- ▶ Microscopic BHs: none
- ▶ Primordial BHs: none (upper bound)
- ▶ Stellar mass BHs in binary systems: many (17 good candidates (including Cygnus X-1), 37 other candidates)
- ▶ Isolated stellar mass BHs: some (1 good candidate, 3 other candidates)
- ▶ Intermediate mass BHs: some (11 candidates)
- ▶ Galactic core BHs: many (Milky Way, 66 other candidates)

Data compiled in 2004 by R. Johnston

Some Black Hole Observational Data

OJ287, about 18 billion solar masses



Artistic impression (NASA Outreach), presented at the Annual meeting of the American Astronomical Society, 2008

- ▶ Microscopic BHs: none
- ▶ Primordial BHs: none (upper bound)
- ▶ Stellar mass BHs in binary systems: many (17 good candidates (including Cygnus X-1), 37 other candidates)
- ▶ Isolated stellar mass BHs: some (1 good candidate, 3 other candidates)
- ▶ Intermediate mass BHs: some (11 candidates)
- ▶ Galactic core BHs: many (Milky Way, 66 other candidates)

Data compiled in 2004 by R. Johnston

Black holes are the simplest explanation of data! Thus, by Occam's razor they exist.

Outline

Black Holes in Gravitational Physics

Black Holes in Non-Gravitational Physics

Case Study: Holographic Renormalization

Black Holes in Science Fiction

All I am going to say about this topic is:

Black Holes in Science Fiction


All I am going to say about this topic is:



Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part I: Picture

Hydraulic jump as a white hole analog



Picture by Piotr Pieranski, taken from a paper by G. Volovik

Some literature:

- ▶ C. Barcelo, S. Liberati,
M. Visser,
[gr-qc/0505065](https://arxiv.org/abs/gr-qc/0505065)
- ▶ G. Volovik,
[gr-qc/0612134](https://arxiv.org/abs/gr-qc/0612134)
- ▶ T. Philbin et al,
[arXiv:0711.4796](https://arxiv.org/abs/0711.4796)
- ▶ M. Visser,
S. Weinfurtner,
[arXiv:0712.0427](https://arxiv.org/abs/0712.0427)

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part II: Formulas

Idea: Linearize perturbations in continuity equation

$$\partial_t \rho + \nabla \cdot (\rho v) = 0$$

and Euler equation

$$\rho(\partial_t v + (v \cdot \nabla)v) = -\nabla p$$

and assume no vorticity, $v = -\nabla\phi$, and barotropic equation of state

$$\nabla h = \frac{1}{\rho} \nabla p$$

Then the velocity-potential ϕ obeys the relativistic (!) wave-equation

$$\square\phi = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \phi) = 0$$

with the acoustic metric

$$g_{\mu\nu}(t, x) = \frac{\rho}{c} \begin{pmatrix} -(c^2 - v^2) & -v^T \\ -v & \mathbb{I} \end{pmatrix}$$

where the speed of sound is given by $c^{-2} = \partial\rho/\partial p$.

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...

...they do not necessarily teach us anything new.



Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...



...they do not necessarily teach us anything new.

Summary

- ▶ Black hole analogs are very useful for pedagogic demonstrations (high schools and undergraduate curricula!)

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...



...they do not necessarily teach us anything new.

Summary

- ▶ Black hole analogs are very useful for pedagogic demonstrations (high schools and undergraduate curricula!)
- ▶ Workers in the field frequently express the hope to experimentally establish the Hawking effect

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...



...they do not necessarily teach us anything new.

Summary

- ▶ Black hole analogs are very useful for pedagogic demonstrations (high schools and undergraduate curricula!)
- ▶ Workers in the field frequently express the hope to experimentally establish the Hawking effect
- ▶ Even if this works it is not clear what it would teach us

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...



...they do not necessarily teach us anything new.

Summary

- ▶ Black hole analogs are very useful for pedagogic demonstrations (high schools and undergraduate curricula!)
- ▶ Workers in the field frequently express the hope to experimentally establish the Hawking effect
- ▶ Even if this works it is not clear what it would teach us
- ▶ But it would be a cool experiment!

Condensed Matter Analogs

Deaf and Dumb holes (W. Unruh 1981), Part III: Reality Check

Black hole analogs are nice, but...



...they do not necessarily teach us anything new.


Summary

- ▶ Black hole analogs are very useful for pedagogic demonstrations (high schools and undergraduate curricula!)
- ▶ Workers in the field frequently express the hope to experimentally establish the Hawking effect
- ▶ Even if this works it is not clear what it would teach us
- ▶ But it would be a cool experiment!

Q: Are there other unexpected applications of black holes?

AdS/CFT Correspondence!

Relates strings on AdS to specific gauge theories at the boundary of AdS



Open Universe Looking from inside, boundary at infinity
Limit Circle IV, by M. C. Escher

Applications

- ▶ RHIC physics
- ▶ Black hole physics
- ▶ Scattering at strong coupling
- ▶ Cold atoms
- ▶ Superconductors
- ▶ Quantum gravity
- ▶ ... to be discovered!

Note: J. Maldacena's paper hep-th/9711200 second most cited paper ever (SPIRES). First is Steven Weinberg's "A Model of Leptons".

Outline

Black Holes in Gravitational Physics

Black Holes in Non-Gravitational Physics

Case Study: Holographic Renormalization

AdS₂

... the simplest gravity model where the need for holographic renormalization arises!

Bulk action:

$$I_B = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right]$$

AdS₂

... the simplest gravity model where the need for holographic renormalization arises!

Bulk action:

$$I_B = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right]$$

Variation with respect to scalar field X yields

$$R = -\frac{2}{\ell^2}$$

This means curvature is constant and negative, i.e., AdS₂.

AdS₂

... the simplest gravity model where the need for holographic renormalization arises!

Bulk action:

$$I_B = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right]$$

Variation with respect to scalar field X yields

$$R = -\frac{2}{\ell^2}$$

This means curvature is constant and negative, i.e., AdS₂.

Variation with respect to metric g yields

$$\nabla_{\mu} \nabla_{\nu} X - g_{\mu\nu} \square X + g_{\mu\nu} \frac{X}{\ell^2} = 0$$

AdS₂

... the simplest gravity model where the need for holographic renormalization arises!

Bulk action:

$$I_B = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right]$$

Variation with respect to scalar field X yields

$$R = -\frac{2}{\ell^2}$$

This means curvature is constant and negative, i.e., AdS₂.

Variation with respect to metric g yields

$$\nabla_{\mu} \nabla_{\nu} X - g_{\mu\nu} \square X + g_{\mu\nu} \frac{X}{\ell^2} = 0$$

Equations of motion above solved by

$$X = r, \quad g_{\mu\nu} dx^{\mu} dx^{\nu} = \left(\frac{r^2}{\ell^2} - M \right) dt^2 + \frac{dr^2}{\frac{r^2}{\ell^2} - M}$$

AdS₂

... the simplest gravity model where the need for holographic renormalization arises!

Bulk action:

$$I_B = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right]$$

Variation with respect to scalar field X yields

$$R = -\frac{2}{\ell^2}$$

This means curvature is constant and negative, i.e., AdS₂.

Variation with respect to metric g yields

$$\nabla_{\mu} \nabla_{\nu} X - g_{\mu\nu} \square X + g_{\mu\nu} \frac{X}{\ell^2} = 0$$

Equations of motion above solved by

$$X = r, \quad g_{\mu\nu} dx^{\mu} dx^{\nu} = \left(\frac{r^2}{\ell^2} - M \right) dt^2 + \frac{dr^2}{\frac{r^2}{\ell^2} - M}$$

There is an important catch, however: Boundary terms tricky!

Boundary terms, Part I

Gibbons–Hawking–York boundary terms: quantum mechanical toy model

Let us start with an **bulk Hamiltonian action**

$$I_B = \int^{t_f} dt [-\dot{p}q - H(q, p)]$$

Boundary terms, Part I

Gibbons–Hawking–York boundary terms: quantum mechanical toy model

Let us start with an bulk Hamiltonian action

$$I_B = \int^{t_f} dt [-\dot{p}q - H(q, p)]$$

Want to set up a Dirichlet boundary value problem $q = \text{fixed}$ at t_f

Boundary terms, Part I

Gibbons–Hawking–York boundary terms: quantum mechanical toy model

Let us start with an bulk Hamiltonian action

$$I_B = \int^{t_f} dt [-\dot{p}q - H(q, p)]$$

Want to set up a Dirichlet boundary value problem $q = \text{fixed}$ at t_f
Problem:

$$\delta I_B = 0 \text{ requires } q \delta p = 0 \text{ at boundary}$$

Boundary terms, Part I

Gibbons–Hawking–York boundary terms: quantum mechanical toy model

Let us start with an **bulk Hamiltonian action**

$$I_B = \int^{t_f} dt [-\dot{p}q - H(q, p)]$$

Want to set up a Dirichlet boundary value problem $q = \text{fixed}$ at t_f
Problem:

$$\delta I_B = 0 \text{ requires } q \delta p = 0 \text{ at boundary}$$

Solution: add “**Gibbons–Hawking–York**” boundary term

$$I_E = I_B + I_{GHY}, \quad I_{GHY} = pq \Big|^{t_f}$$

Boundary terms, Part I

Gibbons–Hawking–York boundary terms: quantum mechanical toy model

Let us start with an bulk Hamiltonian action

$$I_B = \int^{t_f} dt [-\dot{p}q - H(q, p)]$$

Want to set up a Dirichlet boundary value problem $q = \text{fixed}$ at t_f
Problem:

$$\delta I_B = 0 \text{ requires } q \delta p = 0 \text{ at boundary}$$

Solution: add “Gibbons–Hawking–York” boundary term

$$I_E = I_B + I_{GHY}, \quad I_{GHY} = pq \Big|^{t_f}$$

As expected $I_E = \int^{t_f} dt [p\dot{q} - H(q, p)]$ is standard Hamiltonian action

Boundary terms, Part II

Gibbons–Hawking–York boundary terms in gravity — something still missing!

That was easy! In gravity the result is

$$I_{GHY} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K$$

where γ (K) is determinant (trace) of first (second) fundamental form.
Euclidean action with correct boundary value problem is

$$I_E = I_B + I_{GHY}$$

The boundary lies at $r = r_0$, with $r_0 \rightarrow \infty$. Are we done?

Boundary terms, Part II

Gibbons–Hawking–York boundary terms in gravity — something still missing!

That was easy! In gravity the result is

$$I_{GHY} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K$$

where γ (K) is determinant (trace) of first (second) fundamental form.
Euclidean action with correct boundary value problem is

$$I_E = I_B + I_{GHY}$$

The boundary lies at $r = r_0$, with $r_0 \rightarrow \infty$. Are we done?
No! Serious Problem! Variation of I_E yields

$$\delta I_E \sim \text{EOM} + \delta X (\text{boundary} - \text{term}) - \lim_{r \rightarrow \infty} \int_{\partial\mathcal{M}} dt \delta\gamma$$

Boundary terms, Part II

Gibbons–Hawking–York boundary terms in gravity — something still missing!

That was easy! In gravity the result is

$$I_{GHY} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K$$

where γ (K) is determinant (trace) of first (second) fundamental form.
Euclidean action with correct boundary value problem is

$$I_E = I_B + I_{GHY}$$

The boundary lies at $r = r_0$, with $r_0 \rightarrow \infty$. Are we done?
No! Serious Problem! Variation of I_E yields

$$\delta I_E \sim \text{EOM} + \delta X (\text{boundary} - \text{term}) - \lim_{r \rightarrow \infty} \int_{\partial\mathcal{M}} dt \delta\gamma$$

Asymptotic metric: $\gamma = r^2/\ell^2 + \mathcal{O}(1)$. Thus, $\delta\gamma$ may be *finite!*

Boundary terms, Part II

Gibbons–Hawking–York boundary terms in gravity — something still missing!

That was easy! In gravity the result is

$$I_{GHY} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K$$

where γ (K) is determinant (trace) of first (second) fundamental form.
Euclidean action with correct boundary value problem is

$$I_E = I_B + I_{GHY}$$

The boundary lies at $r = r_0$, with $r_0 \rightarrow \infty$. Are we done?
No! Serious Problem! Variation of I_E yields

$$\delta I_E \sim \text{EOM} + \delta X (\text{boundary} - \text{term}) - \lim_{r \rightarrow \infty} \int_{\partial\mathcal{M}} dt \delta\gamma$$

Asymptotic metric: $\gamma = r^2/\ell^2 + \mathcal{O}(1)$. Thus, $\delta\gamma$ may be *finite!*

$\delta I_E \neq 0$ for some variations that preserve boundary conditions!!!

Boundary terms, Part III

Holographic renormalization: quantum mechanical toy model

Key observation: Dirichlet boundary problem not changed under

$$I_E \rightarrow \Gamma = I_E - I_{CT} = I_{EH} + I_{GHY} - I_{CT}$$

with

$$I_{CT} = S(q, t)|^{t_f}$$

Boundary terms, Part III

Holographic renormalization: quantum mechanical toy model

Key observation: Dirichlet boundary problem not changed under

$$I_E \rightarrow \Gamma = I_E - I_{CT} = I_{EH} + I_{GHY} - I_{CT}$$

with

$$I_{CT} = S(q, t)|^{t_f}$$

Improved action:

$$\Gamma = \int^{t_f} dt [-\dot{p}q - H(q, p)] + pq|^{t_f} - S(q, t)|^{t_f}$$

Boundary terms, Part III

Holographic renormalization: quantum mechanical toy model

Key observation: Dirichlet boundary problem not changed under

$$I_E \rightarrow \Gamma = I_E - I_{CT} = I_{EH} + I_{GHY} - I_{CT}$$

with

$$I_{CT} = S(q, t)|^{t_f}$$

Improved action:

$$\Gamma = \int^{t_f} dt [-\dot{p}q - H(q, p)] + pq|^{t_f} - S(q, t)|^{t_f}$$

First variation:

$$\delta\Gamma\Big|_{\text{EOM}} = \left(p - \frac{\partial S(q, t)}{\partial q} \right) \delta q \Big|^{t_f} = 0?$$

Boundary terms, Part III

Holographic renormalization: quantum mechanical toy model

Key observation: Dirichlet boundary problem not changed under

$$I_E \rightarrow \Gamma = I_E - I_{CT} = I_{EH} + I_{GHY} - I_{CT}$$

with

$$I_{CT} = S(q, t)|^{t_f}$$

Improved action:

$$\Gamma = \int^{t_f} dt [-\dot{p}q - H(q, p)] + pq|^{t_f} - S(q, t)|^{t_f}$$

First variation:

$$\delta\Gamma|_{\text{EOM}} = \left(p - \frac{\partial S(q, t)}{\partial q} \right) \delta q|^{t_f} = 0?$$

Works if $S(q, t)$ is Hamilton's principal function!

Boundary terms, Part IV

Holographic renormalization in AdS_2 gravity

Hamilton's principle function

- ▶ Solves the Hamilton–Jacobi equation
- ▶ Does not change boundary value problem when added to action
- ▶ Is capable to render $\delta\Gamma = 0$ even when $\delta I_E \neq 0$
- ▶ Reasonable Ansatz: Holographic counterterm = Solution of Hamilton–Jacobi equation!

Boundary terms, Part IV

Holographic renormalization in AdS₂ gravity

Hamilton's principle function

- ▶ Solves the Hamilton–Jacobi equation
- ▶ Does not change boundary value problem when added to action
- ▶ Is capable to render $\delta\Gamma = 0$ even when $\delta I_E \neq 0$
- ▶ Reasonable Ansatz: Holographic counterterm = Solution of Hamilton–Jacobi equation!

In case of AdS₂ gravity this Ansatz yields

$$I_{\text{CT}} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \frac{X}{\ell}$$

Boundary terms, Part IV

Holographic renormalization in AdS₂ gravity

Hamilton's principle function

- ▶ Solves the Hamilton–Jacobi equation
- ▶ Does not change boundary value problem when added to action
- ▶ Is capable to render $\delta\Gamma = 0$ even when $\delta I_E \neq 0$
- ▶ Reasonable Ansatz: Holographic counterterm = Solution of Hamilton–Jacobi equation!

In case of AdS₂ gravity this Ansatz yields

$$I_{\text{CT}} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \frac{X}{\ell}$$

Action consistent with boundary value problem and variational principle:

$$\Gamma = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right] - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K + \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \frac{X}{\ell}$$

Boundary terms, Part IV

Holographic renormalization in AdS₂ gravity

Hamilton's principle function

- ▶ Solves the Hamilton–Jacobi equation
- ▶ Does not change boundary value problem when added to action
- ▶ Is capable to render $\delta\Gamma = 0$ even when $\delta I_E \neq 0$
- ▶ Reasonable Ansatz: Holographic counterterm = Solution of Hamilton–Jacobi equation!

In case of AdS₂ gravity this Ansatz yields

$$I_{\text{CT}} = - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \frac{X}{\ell}$$

Action consistent with boundary value problem and variational principle:

$$\Gamma = -\frac{1}{2} \int_{\mathcal{M}} d^2x \sqrt{g} \left[X \left(R + \frac{2}{\ell^2} \right) \right] - \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K + \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \frac{X}{\ell}$$

$\delta\Gamma = 0$ for all variations that preserve the boundary conditions!

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The leading term is the 'on-shell' action.

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The **leading term** is the 'on-shell' action.
- ▶ The **linear term** should vanish on solutions g_{cl} and X_{cl} .

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The **leading term** is the 'on-shell' action.
- ▶ The **linear term** should vanish on solutions g_{cl} and X_{cl} .

If nothing goes wrong get partition function

$$\mathcal{Z} \sim \exp\left(-I_E[g_{cl}, X_{cl}]\right) \times \dots$$

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The **leading term** is the 'on-shell' action.
- ▶ The **linear term** should vanish on solutions g_{cl} and X_{cl} .

If nothing goes wrong get partition function

$$\mathcal{Z} \sim \exp\left(-I_E[g_{cl}, X_{cl}]\right) \times \dots$$

Accessibility of the semi-classical approximation requires

1. $I_E[g_{cl}, X_{cl}] > -\infty$
2. $\delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] = 0$

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The **leading term** is the 'on-shell' action.
- ▶ The **linear term** should vanish on solutions g_{cl} and X_{cl} .

If nothing goes wrong get partition function

$$\mathcal{Z} \sim \exp\left(-I_E[g_{cl}, X_{cl}]\right) \times \dots$$

Accessibility of the semi-classical approximation requires

1. $I_E[g_{cl}, X_{cl}] \rightarrow -\infty \rightarrow$ violated in AdS gravity!
2. $\delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] = 0$

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The **leading term** is the 'on-shell' action.
- ▶ The **linear term** should vanish on solutions g_{cl} and X_{cl} .

If nothing goes wrong get partition function

$$\mathcal{Z} \sim \exp\left(-I_E[g_{cl}, X_{cl}]\right) \times \dots$$

Accessibility of the semi-classical approximation requires

1. $I_E[g_{cl}, X_{cl}] \rightarrow -\infty \rightarrow$ violated in AdS gravity!
2. $\delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \neq 0 \rightarrow$ violated in AdS gravity!

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E + \dots$$

- ▶ The **leading term** is the 'on-shell' action.
- ▶ The **linear term** should vanish on solutions g_{cl} and X_{cl} .

If **nothing goes wrong** get partition function

$$\mathcal{Z} \sim \exp\left(-I_E[g_{cl}, X_{cl}]\right) \times \dots$$

Accessibility of the semi-classical approximation requires

1. $I_E[g_{cl}, X_{cl}] \rightarrow -\infty \rightarrow$ violated in AdS gravity!
2. $\delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \neq 0 \rightarrow$ violated in AdS gravity!

Everything goes wrong with I_E !

In particular, do not get correct free energy $F = T I_E = -\infty$ or entropy

$$S = \infty$$

Thermodynamics of Black Holes as a Simple Application

Consider small perturbation around classical solution

$$\Gamma[g_{cl} + \delta g, X_{cl} + \delta X] = \Gamma[g_{cl}, X_{cl}] + \delta\Gamma + \dots$$

- ▶ The leading term is the 'on-shell' action.
- ▶ The linear term should vanish on solutions g_{cl} and X_{cl} .

If nothing goes wrong get partition function

$$\mathcal{Z} \sim \exp\left(-\Gamma[g_{cl}, X_{cl}]\right) \times \dots$$

Accessibility of the semi-classical approximation requires

1. $\Gamma[g_{cl}, X_{cl}] > -\infty \rightarrow$ ok in AdS gravity!
2. $\delta\Gamma[g_{cl}, X_{cl}; \delta g, \delta X] = 0 \rightarrow$ ok in AdS gravity!

Everything works with Γ !

In particular, do get correct free energy $F = TI_E = M - TS$ and entropy

$$S = 2\pi X|_{\text{horizon}} = \text{Area}/4$$

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}
- ▶ Check consistency of variational principle

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}
- ▶ Check consistency of variational principle
- ▶ If necessary, subtract holographic counterterm I_{CT}

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}
- ▶ Check consistency of variational principle
- ▶ If necessary, subtract holographic counterterm I_{CT}
- ▶ Use improved action

$$\Gamma = I_B + I_{GHY} - I_{CT}$$

for applications!

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}
- ▶ Check consistency of variational principle
- ▶ If necessary, subtract holographic counterterm I_{CT}
- ▶ Use improved action

$$\Gamma = I_B + I_{GHY} - I_{CT}$$

for applications!

- ▶ Applications include thermodynamics from Euclidean path integral and calculation of holographic stress tensor in AdS/CFT

$$\delta\Gamma|_{\text{EOM}} \sim \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} T_{ab} \delta\gamma^{ab} \quad \leftrightarrow \quad T_{ab} = \langle T_{ab} \rangle$$

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}
- ▶ Check consistency of variational principle
- ▶ If necessary, subtract holographic counterterm I_{CT}
- ▶ Use improved action

$$\Gamma = I_B + I_{GHY} - I_{CT}$$

for applications!

- ▶ Applications include thermodynamics from Euclidean path integral and calculation of holographic stress tensor in AdS/CFT

$$\delta\Gamma|_{\text{EOM}} \sim \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} T_{ab} \delta\gamma^{ab} \quad \leftrightarrow \quad T_{ab} = \langle T_{ab} \rangle$$

- ▶ Straightforward applications in quantum field theory?

Summary and algorithm of holographic renormalization

In any dimension, for any asymptotics — may arise also in quantum field theory!

- ▶ Start with bulk action I_B
- ▶ Check consistency of boundary value problem
- ▶ If necessary, add boundary term I_{GHY}
- ▶ Check consistency of variational principle
- ▶ If necessary, subtract holographic counterterm I_{CT}
- ▶ Use improved action

$$\Gamma = I_B + I_{GHY} - I_{CT}$$

for applications!






- ▶ Applications include thermodynamics from Euclidean path integral and calculation of holographic stress tensor in AdS/CFT

$$\delta\Gamma|_{\text{EOM}} \sim \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} T_{ab} \delta\gamma^{ab} \quad \leftrightarrow \quad T_{ab} = \langle T_{ab} \rangle$$

- ▶ Straightforward applications in quantum field theory? Possibly!

... and thanks to my collaborators!

Some literature on AdS_2 holography

-  D. Grumiller and R. McNees, JHEP **0704**, 074 [[arXiv:hep-th/0703230](#)].
-  T. Hartman and A. Strominger, “Central charge for AdS_2 quantum gravity,” [[arXiv:0803.3621 \[hep-th\]](#)].
-  M. Alishahiha and F. Ardalan, “Central charge for 2D gravity on AdS_2 and $\text{AdS}_2/\text{CFT}_1$ correspondence,” [[arXiv:0805.1861 \[hep-th\]](#)].
-  R. Gupta and A. Sen, “ $\text{AdS}(3)/\text{CFT}(2)$ to $\text{AdS}(2)/\text{CFT}(1)$,” [[arXiv:0806.0053 \[hep-th\]](#)].
-  A. Castro, D. Grumiller, R. McNees and F. Larsen, “Holographic description of AdS_2 black holes,” [[arXiv:0809.4264 \[hep-th\]](#)].

Thank you for your attention!